

ADDENDUM: ON THE TYPE OF THE SPIN POLARIZATION DEPENDENCE OF THE NEUTRINO MASS AND CHARGE

Rasulkhozha S. Sharafiddinov

Institute of Nuclear Physics, Uzbekistan Academy of Sciences,
Tashkent, 702132 Ulugbek, Uzbekistan

We discuss the characteristic features of the latent nature of the neutrino mass, according to which, all components of leptonic current can appear in the interaction type dependence. Such a regularity, however, requires the modification of the some denotations in the corresponding formulas of the paper *On the type of the spin polarization dependence of the neutrino mass and charge* [1]. We will also include in a given letter the full version of the original article with necessary replacements of the structural variables. They of course do not change our implications.

In our original paper [1] the question has been raised as to whether there exists any spin polarization type dependence of the neutrino mass, charge and behavior. This sight at the nature of the neutrino is based logically on the analysis of the possible processes of elastic scattering of longitudinal and transversal polarized light neutrinos ($\nu = \nu_e$) by spinless nuclei in the presence of the Coulomb m_ν^E , weak m_ν^W and the united electroweak m_ν^{EW} masses, and also the charge $f_{1\nu}$, magnetic $f_{2\nu}$, anapole $g_{1\nu}$ and electric dipole $g_{2\nu}$ moments of incoming fermions with neutral currents. At the definition of their cross sections, we used the matrix elements of transitions in the form

$$M_{fi}^E = \frac{4\pi\alpha}{q_E^2} \bar{u}(p'_E, s') [\gamma_\mu f_{1\nu}(0) - i\sigma_{\mu\lambda} q_E^\lambda f_{2\nu}(0) + \gamma_5 \gamma_\mu g_{1\nu}(0) - i\gamma_5 \sigma_{\mu\lambda} q_E^\lambda g_{2\nu}(0)] u(p_E, s) J_\mu^\gamma(q_E), \quad (1)$$

$$M_{fi}^W = \frac{G_F}{\sqrt{2}} \bar{u}(p'_W, s') \gamma_\mu (g_{V_\nu} + \gamma_5 g_{A_\nu}) u(p_W, s) J_\mu^{Z^0}(q_W), \quad (2)$$

where $q_E = p_E - p'_E$, $q_W = p_W - p'_W$, $p_E(p_W)$ and $p'_E(p'_W)$ denote the four-momentum of a particle before and after the electric (weak) scattering, s and s' are the helicities of these states, J_μ^γ and $J_\mu^{Z^0}$ are the nuclear currents in the studied processes, g_{V_ν} and g_{A_ν} correspond to the vector and axial-vector components of leptonic weak neutral currents.

According to our description, the electric mass of the neutrino leads to its Coulomb scattering. At the exchange by an Z^0 -boson, only the weak component of mass is responsible for the scattering.

In the framework of mass-charge duality [2], each of the existing types of charges testifies in favor of the availability of a kind of inertial mass. Any particle of a Dirac nature can, therefore, have simultaneously both electric and weak [3] masses.

Thus, in the mentioned paper [1] the neutrino scattering by nuclei of an electroweak charge going at the expense of the interference between the interactions (1) and (2) is analytically described as

$$2\text{Re} M_{fi}^E M_{fi}^{*W} = \frac{8\pi\alpha G_F}{\sqrt{2} q_{EW}^2} \text{Re} \rho_{EW} \rho'_{EW} [\gamma_\mu f_{1\nu}(0) - i\sigma_{\mu\lambda} q_{EW}^\lambda f_{2\nu}(0) + \gamma_5 \gamma_\mu g_{1\nu}(0) - i\gamma_5 \sigma_{\mu\lambda} q_{EW}^\lambda g_{2\nu}(0)] \gamma_\mu (g_{V_\nu} + \gamma_5 g_{A_\nu}) J_\mu^\gamma(q_{EW}) J_\mu^{Z^0}(q_{EW}) \quad (3)$$

and it is assumed additionally that

$$q_{EW} = p_{EW} - p'_{EW}, \quad m_\nu^{EW} = m_\nu^E + m_\nu^W,$$

$$\rho_{EW} = u(p_{EW}, s) \bar{u}(p_{EW}, s), \quad \rho'_{EW} = u(p'_{EW}, s') \bar{u}(p'_{EW}, s').$$

On the other hand, as follows from considerations of symmetry, any component of leptonic current there exists at the availability of a kind of charge [4]. If this is so, each of the interactions (1) and (3) states that the terms $f_{i\nu}$ and $g_{i\nu}$ responsible for the interference scattering are not of those form factors which arise as a result of a particle Coulomb mass. The account of their differences leads (1) to the following:

$$M_{fi}^E = \frac{4\pi\alpha}{q_E^2} \bar{u}(p'_E, s') [\gamma_\mu f_{1\nu}^E(0) - i\sigma_{\mu\lambda} q_E^\lambda f_{2\nu}^E(0) +$$

$$+ \gamma_5 \gamma_\mu g_{1\nu}^E(0) - i\gamma_5 \sigma_{\mu\lambda} q_E^\lambda g_{2\nu}^E(0)] u(p_E, s) J_\mu^\gamma(q_E) \quad (4)$$

and that, consequently, the interactions (3) must have the form

$$2\text{Re} M_{fi}^E M_{fi}^{*W} = \frac{8\pi\alpha G_F}{\sqrt{2}q_{EW}^2} \text{Re} \Lambda_{EW} \Lambda'_{EW} [\gamma_\mu f_{1\nu}^I(0) -$$

$$- i\sigma_{\mu\lambda} q_{EW}^\lambda f_{2\nu}^I(0) + \gamma_5 \gamma_\mu g_{1\nu}^I(0) -$$

$$- i\gamma_5 \sigma_{\mu\lambda} q_{EW}^\lambda g_{2\nu}^I(0)] \gamma_\mu (g_{V\nu} + \gamma_5 g_{A\nu}) J_\mu^\gamma(q_{EW}) J_\mu^{Z^0}(q_{EW}). \quad (5)$$

Here $\Lambda_{EW} = \rho_{EW}$ and $\Lambda'_{EW} = \rho'_{EW}$. Unlike $f_{i\nu}^E$ and $g_{i\nu}^E$, the interference terms $f_{i\nu}^I$ and $g_{i\nu}^I$ characterize the united electroweak mass dependence of the neutrino form factors.

Inserting (2), (4) and (5) in

$$\frac{d\sigma_{E,W}(s, s')}{d\Omega} = \frac{1}{16\pi^2} |M_{fi}^E + M_{fi}^W|^2, \quad (6)$$

we find for the elastic scattering cross sections an explicit expressions which coincide with the corresponding formulas from the paper [1] if make there the following replacements:

$$f_{i\nu} \rightarrow f_{i\nu}^t, \quad g_{i\nu} \rightarrow g_{i\nu}^t,$$

$$F_{EW}(q_{EW}^2) \rightarrow F_I(q_{EW}^2), \quad \theta \rightarrow \theta_K,$$

$$\sigma_o^\nu \rightarrow \sigma_o^j, \quad \rho \rightarrow \rho_{EW},$$

$$t = E, I, \quad j = E, EW, \quad K = E, EW, W.$$

Of them θ_K denotes the scattering angle in the mass type dependence, F_I is the nuclear interference form factor, σ_o^j describes the Mott cross section at the neutrino interaction in the presence of one of the masses, m_ν^E or m_ν^{EW} , and ρ_{EW} is the electroweak interference coefficient.

Thus, the general picture is obtained of elastic scattering in which an important part is said of the compound structure of mass and charge.

Therefore, we will include in this letter the full version of the paper [1] with the above noted modifications. They of course do not change our implications based on the analysis of earlier findings.

References

- [1] R.S. Sharafiddinov, Spacetime Subst. **3**, 134 (2002); physics/0305015, v 2.
- [2] R.S. Sharafiddinov, Spacetime Subst. **3**, 47 (2002); physics/0305008.
- [3] S. Weinberg, Phys. Rev. Lett. **29**, 388 (1972).
- [4] R.S. Sharafiddinov, Spacetime Subst. **3**, 86 (2002); physics/0305009.

ON THE TYPE OF THE SPIN POLARIZATION DEPENDENCE OF THE NEUTRINO MASS AND CHARGE

Rasulkhozha S. Sharafiddinov

**Institute of Nuclear Physics, Uzbekistan Academy of Sciences,
Tashkent, 702132 Ulugbek, Uzbekistan**

Any non-zero component of charge implies the existence of a kind of inertial mass. Therefore, each of existing types of the dipole moments must arise as a consequence of the availability of a kind of charge. From their point of view, the elastic scattering of completely longitudinally (transversally) polarized neutrinos and antineutrinos by spinless nuclei is discussed taking into account the Coulomb, weak and the united electroweak masses and charges, and also the magnetic, anapole and electric dipole moments of incoming fermions with neutral currents. Interconversions of neutrinos of the different components have been investigated, at which a particle chiral invariance is violated at the expense of the flip of its spin. This becomes possible owing to an intimate connection between these phenomena and character of the structure of a Dirac mass. Analysis of the studied process cross sections assumed that both masses and charges of longitudinal and transversal neutrinos are strictly unidentical.

1. Introduction

In studying the nature of elementary particles such characteristics as the mass and charge play a large role. At the same time it is well known that according to the hypothesis of field mass based on the classical theory of an extensive electron [1], a particle all the mass is purely electric. Such a structure, however, meets with many problems. One of them states that the charge distribution of the electron is not steady.

My investigation of the interaction of electrons and their neutrinos with field of emission shows clearly [2, 3] that if the neutrino corresponds to the electron ($\nu = \nu_e$), its full electric charge e_ν^{full} and magnetic moment μ_ν^{full} appear owing to the Coulomb mass m_ν^E and behave as [4]

$$e_\nu^{full} = -\frac{3eG_F m_\nu^2}{4\pi^2 \sqrt{2}}, \quad e = |e|, \quad (1)$$

$$\mu_\nu^{full} = \frac{3eG_F m_\nu}{8\pi^2 \sqrt{2}}, \quad m_\nu = m_\nu^E. \quad (2)$$

This picture reflects the fact that each of the available types of charges says in favor of the availability of a kind of inertial mass. Thereby the possibility of the existence of the united rest mass m_ν^U and charge e_ν^U for the neutrino equal, respectively, to its all the mass and charge is not excluded. One can define their structure in the form [5]

$$m_\nu^U = m_\nu^E + m_\nu^W + m_\nu^S + \dots, \quad (3)$$

$$e_\nu^U = e_\nu^E + e_\nu^W + e_\nu^S + \dots \quad (4)$$

Here the indices E , W and S correspond to the electric, weak and strong components of the neutrino mass and charge. They constitute herewith the harmony of forces of the different nature. Therefore, the charge distributions of light leptons must be steady [6].

Such a stability of matter saying about the compound structure of charge quantization law [7] and testifies of that any non-zero component of the electric charge implies the existence of a kind of the dipole moment. In other words, the Dirac $F_{1\nu}(q^2)$ and the Pauli $F_{2\nu}(q^2)$ form factors contain the static as well as the dynamical components [6]:

$$F_{i\nu}(q^2) = f_{i\nu}(0) + A_{i\nu}(\vec{q}^2) + \dots, \quad (5)$$

where $f_{i\nu}(0)$ give the normal size of charge and moment, $A_{i\nu}(\vec{q}^2)$ characterize the momentum dependence of leptonic current vector parts. The terms $f_{i\nu}(0)$ and $A_{i\nu}(\vec{q}^2)$ are responsible for the first and the second Born approximations. In these circumstances, the form factors $F_{1\nu}(q^2)$ and $F_{2\nu}(q^2)$ at $q^2 = 0$ define the full static values of the neutrino electric charge and moment:

$$F_{1\nu}(0) = e_\nu^{full} = e_\nu^{norm} + e_\nu^{anom} + \dots, \quad (6)$$

$$F_{2\nu}(0) = \mu_\nu^{full} = \mu_\nu^{norm} + \mu_\nu^{anom} + \dots \quad (7)$$

Of course, the electric mass and charge of a Dirac neutrino correspond to the most diverse forms of the same regularity of its Coulomb nature. By this reason we conclude that the neutrino electric mass m_ν^E includes as well as the normal m_ν^{norm} and anomalous m_ν^{anom} components [4]:

$$m_\nu^E = m_\nu^{norm} + m_\nu^{anom} + \dots \quad (8)$$

However, it is known [8, 9] that the neutrino interaction with virtual photon is described by the vertex operator

$$\begin{aligned} \Gamma_\mu(p, p') = & \bar{u}(p', s') [\gamma_\mu F_{1\nu}(q^2) - i\sigma_{\mu\lambda} q_\lambda F_{2\nu}(q^2) + \\ & + \gamma_5 \gamma_\mu G_{1\nu}(q^2) - i\gamma_5 \sigma_{\mu\lambda} q_\lambda G_{2\nu}(q^2)] u(p, s). \end{aligned} \quad (9)$$

Here $\sigma_{\mu\lambda} = [\gamma_\mu, \gamma_\lambda]/2$, $q = p - p'$, $p(s)$ and $p'(s')$ denote the four-momentum (helicities) of the neutrino before and after emission, $G_{1\nu}(q^2)$ and $G_{2\nu}(q^2)$ are the leptonic current axial-vector parts.

Analysis of electroweak processes on nuclei assumed [10] that $F_{2\nu}$ and $G_{2\nu}$ must have the same size. Therefore, without loss of generality, we may write

$$G_{i\nu}(q^2) = g_{i\nu}(0) + \Phi_{i\nu}(\vec{q}^2) + \dots, \quad (10)$$

where $g_{1\nu}(0)$ and $g_{2\nu}(0)$ are the normal components of a particle anapole [11] and electric dipole moments. The functions $\Phi_{i\nu}(\vec{q}^2)$ characterize the anomalous behavior of the axial-vector form factors.

According to the correspondence principle, each term in (10) as well as in (5), corresponds to the definite approximation. This sight at the interaction axial-vector nature quality explains the fact that $G_{1\nu}(0)$ and $G_{2\nu}(0)$ give the full static size of a Dirac particle anapole and electric dipole moments:

$$G_{1\nu}(0) = a_\nu^{full} = a_\nu^{norm} + a_\nu^{anom} + \dots, \quad (11)$$

$$G_{2\nu}(0) = d_\nu^{full} = d_\nu^{norm} + d_\nu^{anom} + \dots \quad (12)$$

Of them a_ν^{full} also can be measured experimentally [12]. For d_ν^{full} as well as for e_ν^{full} and μ_ν^{full} there exist laboratory and cosmological restrictions [13, 14].

The purpose of the present work is to discuss the problem of the neutrino latent mass investigating its interaction with field of electroweak emission in the polarization type dependence. First of all we consider the elastic scattering of longitudinal polarized massive Dirac neutrinos by nuclei of the electric (Z) and weak (Z_W) charges

$$\nu_{L,R}(\bar{\nu}_{R,L}) + A(Z, Z_W) \xrightarrow{\gamma, Z^0} \nu'(\bar{\nu}') + A(Z, Z_W) \quad (13)$$

at the account of their Coulomb, weak and the united electroweak masses and currents. Next, all they will be reanalyzed for the transversal case of an incoming particle polarization. In conclusion we present some implications implied from these considerations.

2. Interaction of Longitudinal Polarized Neutrinos with Field of a Nucleus

From our earlier developments, we find that the matrix elements of elastic scattering of an arbitrary polarized neutrinos on the nucleus electric and weak charges in the first Born approximation must have the following structure:

$$M_{fi}^E = \frac{4\pi\alpha}{q_E^2} \bar{u}(p'_E, s') [\gamma_\mu f_{1\nu}^E(0) - i\sigma_{\mu\lambda} q_E^\lambda f_{2\nu}^E(0) + \gamma_5 \gamma_\mu g_{1\nu}^E(0) - i\gamma_5 \sigma_{\mu\lambda} q_E^\lambda g_{2\nu}^E(0)] u(p_E, s) J_\mu^\gamma(q_E), \quad (14)$$

$$M_{fi}^W = \frac{G_F}{\sqrt{2}} \bar{u}(p'_W, s') \gamma_\mu (g_{V\nu} + \gamma_5 g_{A\nu}) u(p_W, s) J_\mu^{Z^0}(q_W). \quad (15)$$

Here $\nu = \nu_{L,R} = \nu_{eL,R}$, $q_E = p_E - p'_E$, $q_W = p_W - p'_W$, $p_E(p_W)$ and $p'_E(p'_W)$ imply the four-momentum of the neutrino before and after the electric (weak) emission, $f_{i\nu}^E$ and $g_{i\nu}^E$ characterize the Coulomb m_ν^E mass dependence of the neutrino form factors, J_μ^x are the nuclear photon ($x = \gamma$) and weak neutral ($x = Z^0$) currents [15], $g_{V\nu}$ and $g_{A\nu}$ denote the corresponding constants of purely weak interaction vector and axial-vector components.

As seen from (14) and (15), in the case of exchange by the photon, only the Coulomb mass is responsible for the electric scattering. Insofar as m_ν^W is concerned, it leads to the corresponding weak interaction.

It appears that here on the basis of the standard definition

$$\frac{d\sigma_{E,W}(s, s')}{d\Omega} = \frac{1}{16\pi^2} |M_{fi}^E + M_{fi}^W|^2 \quad (16)$$

one can will establish an explicit form of the studied process cross sections. It is not excluded, however, that any Dirac particle possesses simultaneously both electric and weak masses. In other words, the neutrino interaction with field of emission explained by an electroweak interference arises at the expense of exchange simultaneously both by the photon and by the weak boson. From this point of view, the interference between the interactions (14) and (15) can be expressed as follows:

$$\begin{aligned} 2\text{Re}M_{fi}^E M_{fi}^{*W} = & \frac{8\pi\alpha G_F}{\sqrt{2}q_{EW}^2} \text{Re}\Lambda_{EW}\Lambda'_{EW} [\gamma_\mu f_{1\nu}^I(0) - \\ & - i\sigma_{\mu\lambda} q_{EW}^\lambda f_{2\nu}^I(0) + \gamma_5 \gamma_\mu g_{1\nu}^I(0) - \\ & - i\gamma_5 \sigma_{\mu\lambda} q_{EW}^\lambda g_{2\nu}^I(0)] \gamma_\mu (g_{V_\nu} + \gamma_5 g_{A_\nu}) J_\mu^\gamma(q_{EW}) J_\mu^{Z^0}(q_{EW}), \end{aligned} \quad (17)$$

where the interaction of each of the currents $f_{i\nu}^I$ and $g_{i\nu}^I$ with the united field of emission of the photon and weak boson is explained by the electroweak structures of mass and charge. They are of course strictly interference. Here and further it is also necessary to keep in mind that

$$q_{EW} = p_{EW} - p'_{EW}, \quad m_\nu^{EW} = m_\nu^E + m_\nu^W,$$

$$\Lambda_{EW} = u(p_{EW}, s) \bar{u}(p_{EW}, s), \quad \Lambda'_{EW} = u(p'_{EW}, s') \bar{u}(p'_{EW}, s').$$

According to these data, the general form of the cross section of the process (13) at the account of longitudinal polarization of both incoming and outgoing fermions is written in the form

$$\begin{aligned} d\sigma_{E,W}^{V,A}(\theta_{E,W}, s, s') = & d\sigma_E^{V,A}(\theta_E, s, s') + \\ & + d\sigma_I^{V,A}(\theta_{EW}, s, s') + d\sigma_W^{V,A}(\theta_W, s, s'), \end{aligned} \quad (18)$$

where the contribution of purely electric mass has the size

$$\begin{aligned} \frac{d\sigma_E^{V,A}(\theta_E, s, s')}{d\Omega} = & \frac{1}{2} \sigma_o^E (1 - \eta_E^2)^{-1} \{ (1 + ss') [f_{1\nu}^E + \\ & + 2\lambda_c s \sqrt{1 - \eta_E^2} g_{1\nu}^E] f_{1\nu}^E + \end{aligned}$$

$$\begin{aligned}
& +\eta_E^2(1-ss')[(f_{1\nu}^E)^2 + 4(m_\nu^E)^2(1-\eta_E^{-2})^2(f_{2\nu}^E)^2]tg^2\frac{\theta_E}{2}- \\
& -8s(E_\nu^E)^2(1-ss')(1-\eta_E^2)^{3/2}f_{2\nu}^E g_{2\nu}^E tg^2\frac{\theta_E}{2}+ \\
& +(1-\eta_E^2)[(1+ss')(g_{1\nu}^E)^2+ \\
& +4(E_\nu^E)^2(1-ss')(g_{2\nu}^E)^2tg^2\frac{\theta_E}{2}]\}F_E^2(q_E^2). \tag{19}
\end{aligned}$$

The second term characterize the interference of the electric and weak interactions as a consequence of the availability of the united electroweak rest mass of incoming particles:

$$\begin{aligned}
\frac{d\sigma_I^{V,A}(\theta_{EW}, s, s')}{d\Omega} &= \frac{1}{2}\rho_{EW}\sigma_o^{EW}(1-\eta_{EW}^2)^{-1}g_{V_\nu}\{(1+ss')[1- \\
& -\lambda_c s\frac{g_{A_\nu}}{g_{V_\nu}}\sqrt{1-\eta_{EW}^2}][f_{1\nu}^I + \lambda_c s\sqrt{1-\eta_{EW}^2}g_{1\nu}^I]+ \\
& +\eta_{EW}^2(1-ss')f_{1\nu}^I tg^2\frac{\theta_{EW}}{2}\}F_I(q_{EW}^2). \tag{20}
\end{aligned}$$

The corresponding cross section for the process going at the expense of the neutrino purely weak rest mass behaves as

$$\begin{aligned}
\frac{d\sigma_W^{V,A}(\theta_W, s, s')}{d\Omega} &= \frac{G_F^2(E_\nu^W)^2}{16\pi^2}\{g_{V_\nu}^2[(1+ss')\cos^2\frac{\theta_W}{2}+ \\
& +\eta_W^2(1-ss')\sin^2\frac{\theta_W}{2}] + g_{A_\nu}^2(1+ss')(1-\eta_W^2)\cos^2\frac{\theta_W}{2}- \\
& -2\lambda_c s g_{V_\nu} g_{A_\nu}(1+ss')\sqrt{1-\eta_W^2}\cos^2\frac{\theta_W}{2}\}F_W^2(q_W^2). \tag{21}
\end{aligned}$$

Here we must have in view of that

$$\begin{aligned}
\sigma_o^E &= \frac{\alpha^2 \cos^2(\theta_E/2)}{4(E_\nu^E)^2(1-\eta_E^2)\sin^4(\theta_E/2)}, \quad \rho_{EW} = \frac{G_F q_{EW}^2}{2\pi\sqrt{2}\alpha}, \\
\sigma_o^{EW} &= \frac{\alpha^2 \cos^2(\theta_{EW}/2)}{4(E_\nu^{EW})^2(1-\eta_{EW}^2)\sin^4(\theta_{EW}/2)}, \quad \eta_E = \frac{m_\nu^E}{E_\nu^E}, \\
\eta_{EW} &= \frac{m_\nu^{EW}}{E_\nu^{EW}}, \quad \eta_W = \frac{m_\nu^W}{E_\nu^W}, \quad F_E(q_E^2) = ZF_c(q_E^2), \\
F_I(q_{EW}^2) &= ZZ_W F_c^2(q_{EW}^2), \quad F_W(q_W^2) = Z_W F_c(q_W^2),
\end{aligned}$$

$$Z_W = \frac{1}{2}\{\beta_V^{(0)}(Z + N) + \beta_V^{(1)}(Z - N)\},$$

$$A = Z + N, \quad M_T = \frac{1}{2}(Z - N),$$

where θ_E , θ_{EW} and θ_W are the polar angles in the electric, electroweak and purely weak scattering at the neutrino energies E_ν^E , E_ν^{EW} and E_ν^W , the functions $F_c(q_E^2)$, $F_c(q_{EW}^2)$ and $F_c(q_W^2)$ describe the charge ($F_c(0) = 1$) form factors of a nucleus in these three processes, M_T is the projection of its isospin T , and constants $\beta_V^{(0)}$ and $\beta_V^{(1)}$ correspond to the isoscalar and isovector components of the nuclear vector neutral current.

The indices V and A imply the simultaneous presence of both vector and axial-vector parts of leptonic photon and weak neutral currents. Therefore, any of the expressions (19)-(21) for the neutrino ($\lambda_c = +1$) and the antineutrino ($\lambda_c = -1$) is different which becomes possible owing to the interference of the interaction vector and axial-vector components.

The terms $(1 + ss')$ and $(1 - ss')$ characterize the scattering with conservation ($s' = s$) and change ($s' = -s$) of helicities of incoming left ($s = -1$) - and right ($s = +1$)-polarized particles. Under such circumstances the value of the cross section (18) conveniently replace by the summed size

$$d\sigma_{E,W}^{V,A}(\theta_{E,W}, s) = d\sigma_{E,W}^{V,A}(\theta_{E,W}, s' = s) + d\sigma_{E,W}^{V,A}(\theta_{E,W}, s' = -s). \quad (22)$$

The compound structures of both terms of (22) testify of that the neutrino charge $f_{1\nu}$ leads to the scattering either with or without flip of its spin. The anapole $g_{1\nu}$ does not change a particle helicity. In contrast to this, the vector $f_{2\nu}$ and axial-vector $g_{2\nu}$ moments are responsible only for the flip of the neutrino spin. Of course, our formulas can also confirm the fact that the helicity of the neutrino of large energy ($E_\nu \gg m_\nu$) is not changed.

However, as known, the right-handed neutrino meets with the problem which states that a chiral symmetry characterized a massive particle does not exist. Therefore, it appears that the neutrinos have no neither the electric, weak nor any other mass. But this is not quite so. The point is that at the helicity conservation a particle chirality is not changed even if it possesses a non-zero rest mass. In other words, the longitudinal neutrino chirality can be violated at the expense of mass, charge, magnetic and electric dipole moments because they lead to the flip of its spin.

In the absence of one of the currents, V or A , any of (19)-(21) not only for the particles and antiparticles but also for the left - and right-handed neutrinos coincides. Such an equality takes place as well as in the low energy limits of the corresponding processes.

3. Transversal Polarized Neutrino Scattering by Nuclei of Electroweak Charges

Owing to an intimate connection between the mass of a particle and its physical nature, any massive neutrino has the longitudinal as well as the transversal polarization. Here an important circumstance is the fact that the same neutrino must not be simultaneously both longitudinal and transversal fermions. There exists, however, the possibility that the longitudinal polarized neutrinos in the elastic scattering on a nucleus can be converted into the transversal polarized and vice versa [16].

Returning to (14)-(17), for the transversal case of the neutrino polarization we establish the compound structure of the cross section of the process (13) which one can present as follows:

$$\begin{aligned} d\sigma_{E,W}^{V,A}(\theta_{E,W}, \varphi, s, s') &= d\sigma_E^{V,A}(\theta_E, \varphi, s, s') + \\ &+ d\sigma_I^{V,A}(\theta_{EW}, \varphi, s, s') + d\sigma_W^{V,A}(\theta_W, \varphi, s, s'). \end{aligned} \quad (23)$$

Here to the contribution of purely electric mass answers the expression

$$\begin{aligned} \frac{d\sigma_E^{V,A}(\theta_E, \varphi, s, s')}{d\Omega} &= \frac{1}{2}\sigma_o^E(1 - \eta_E^2)^{-1}\{((1 + ss')\alpha_T \cos^2 \frac{\varphi}{2} + \\ &+ (1 - ss')\alpha_T^* \sin^2 \frac{\varphi}{2})(f_{1\nu}^E)^2 + \eta_E^2((1 + ss')\gamma_T \sin^2 \frac{\varphi}{2} - \\ &- (1 - ss')\gamma_T^* \cos^2 \frac{\varphi}{2})[(f_{1\nu}^E)^2 + 4(m_\nu^E)^2(1 - \eta_E^{-2})^2(f_{2\nu}^E)^2]tg^2 \frac{\theta_E}{2} + \\ &+ 2\lambda_c s \eta_E \sqrt{1 - \eta_E^2}((1 + ss')\sin^2 \frac{\varphi}{2} - \\ &- (1 - ss')\cos^2 \frac{\varphi}{2})\gamma_T^* f_{1\nu}^E g_{1\nu}^E tg \frac{\theta_E}{2} + \\ &+ (1 - \eta_E^2)((1 + ss')\alpha_T^* \sin^2 \frac{\varphi}{2} + (1 - ss')\alpha_T \cos^2 \frac{\varphi}{2})(g_{1\nu}^E)^2 + \\ &+ 4(E_\nu^E)^2(1 - \eta_E^2)((1 + ss')\gamma_T^* \cos^2 \frac{\varphi}{2} - \end{aligned}$$

$$- (1 - ss')\gamma_T \sin^2 \frac{\varphi}{2} (g_{2\nu}^E)^2 t g^2 \frac{\theta_E}{2} \} F_E^2(q_E^2). \quad (24)$$

The cross section of the interference process originated at the expense of the united electroweak rest mass of incoming transversal polarized neutrinos has the following structure:

$$\begin{aligned} \frac{d\sigma_I^{V,A}(\theta_{EW}, \varphi, s, s')}{d\Omega} = & \frac{1}{2} \rho_{EW} \sigma_o^{EW} (1 - \eta_{EW}^2)^{-1} g_{V_\nu} \{ ((1 + ss')\alpha_T \cos^2 \frac{\varphi}{2} + \\ & + (1 - ss')\alpha_T^* \sin^2 \frac{\varphi}{2}) f_{1\nu}^I + \\ & + \eta_{EW}^2 [((1 + ss')\gamma_T \sin^2 \frac{\varphi}{2} - (1 - ss')\gamma_T^* \cos^2 \frac{\varphi}{2}) t g \frac{\theta_{EW}}{2} + \\ & + \lambda_c s \frac{g_{A_\nu}}{g_{V_\nu}} \eta_{EW}^{-1} \sqrt{1 - \eta_{EW}^2} ((1 + ss') \sin^2 \frac{\varphi}{2} - \\ & - (1 - ss') \cos^2 \frac{\varphi}{2}) \gamma_T^*] f_{1\nu}^I t g \frac{\theta_{EW}}{2} - \\ & - \lambda_c s \eta_{EW} \sqrt{1 - \eta_{EW}^2} [((1 + ss') \sin^2 \frac{\varphi}{2} - \\ & - (1 - ss') \cos^2 \frac{\varphi}{2}) \gamma_T^* t g \frac{\theta_{EW}}{2} + \\ & + \lambda_c s \frac{g_{A_\nu}}{g_{V_\nu}} \eta_{EW}^{-1} \sqrt{1 - \eta_{EW}^2} ((1 + ss') \alpha_T^* \sin^2 \frac{\varphi}{2} + \\ & + (1 - ss') \alpha_T \cos^2 \frac{\varphi}{2})] g_{1\nu}^I \} F_I(q_{EW}^2). \end{aligned} \quad (25)$$

The contribution explained by the transversal polarized neutrinos purely weak rest mass is written in the form

$$\begin{aligned} \frac{d\sigma_W^{V,A}(\theta_W, \varphi, s, s')}{d\Omega} = & \frac{G_F^2 (E_\nu^W)^2}{16\pi^2} \{ g_{V_\nu}^2 [((1 + ss')\alpha_T \cos^2 \frac{\varphi}{2} + \\ & + (1 - ss')\alpha_T^* \sin^2 \frac{\varphi}{2}) c t g^2 \frac{\theta_W}{2} + \\ & + \eta_W^2 ((1 + ss')\gamma_T \sin^2 \frac{\varphi}{2} - (1 - ss')\gamma_T^* \cos^2 \frac{\varphi}{2})] \sin^2 \frac{\theta_W}{2} + \\ & + g_{A_\nu}^2 (1 - \eta_W^2) ((1 + ss')\alpha_T^* \sin^2 \frac{\varphi}{2} + \\ & + (1 - ss')\alpha_T \cos^2 \frac{\varphi}{2}) \cos^2 \frac{\theta_W}{2} - \end{aligned}$$

$$\begin{aligned}
& -2\lambda_c s g_{V_\nu} g_{A_\nu} \eta_W \sqrt{1 - \eta_W^2} ((1 + ss') \sin^2 \frac{\varphi}{2} - \\
& - (1 - ss') \cos^2 \frac{\varphi}{2}) \gamma_T^* \sin \frac{\theta_W}{2} \cos \frac{\theta_W}{2} \} F_W^2(q_W^2),
\end{aligned} \tag{26}$$

where it has been accepted that

$$\begin{aligned}
\alpha_T &= 1 - 2(1 - 4\sin^2 \frac{\varphi}{2}) \sin^2 \frac{\varphi}{2}, \\
\alpha_T^* &= 1 + 2(1 - 4\sin^2 \frac{\varphi}{2}) \cos^2 \frac{\varphi}{2}, \\
\gamma_T &= 1 + 2\cos^2 \frac{\varphi}{2}, \quad \gamma_T^* = 1 - 2\cos^2 \frac{\varphi}{2}.
\end{aligned}$$

Here φ is the azimuthal angle.

As well as in (18), each term in (23) contains the contributions of vector and axial-vector interactions, and also the contributions of their interference between themselves owing to which, the neutrino and antineutrino scattering cross sections are different.

Furthermore, if taken into account the availability of the multipliers $(1 + ss')$ and $(1 - ss')$ in (24)-(26), we can present (23) in the form

$$d\sigma_{E,W}^{V,A}(\theta_{E,W}, \varphi, s) = d\sigma_{E,W}^{V,A}(\theta_{E,W}, \varphi, s' = s) + d\sigma_{E,W}^{V,A}(\theta_{E,W}, \varphi, s' = -s). \tag{27}$$

An explicit expressions for both terms of (27) have the most diverse structures. From their point of view, in the case of the neutrino transversal polarization, the processes with or without change of incoming particle helicities must go as a consequence not only of charge but also of any dipole moment. It is of course not excluded that the flip of the transversal neutrino spin which arises at the expense of mass, charge, magnetic, anapole and electric dipole moments can explain the possible violation of its chirality.

The absence of one of the currents, V or A , implies that each of (24)-(26) for the neutrino and antineutrino as well as for the left - and right-handed particles is not different. Such a coincidence takes place even in the low energy limits of the corresponding types of interactions.

4. Conclusion

In conformity with the laws of nature of a massive neutrino, the presence of any type charge implies the existence of a kind of inertial mass [5]. Such

a duality of matter saying about the steadiness of charge distribution of the neutrino and testifies of that each of all possible types of the dipole moments arises as a consequence of the availability of a kind of charge [6]. Therefore, to reanalyze these features and discuss their some implications we have established the compound structures of the differential cross sections describing the elastic scattering of completely longitudinally (transversally) polarized neutrinos and antineutrinos by spinless nuclei taking into account the Coulomb, weak and the united electroweak masses and charges, and also the magnetic, anapole and electric dipole moments of incoming fermions with neutral currents.

They state that if neutrinos are of longitudinal polarized, their charge answers to the elastic scattering either with or without flip of the spin. The anapole is responsible only for a particle helicity conservation. Unlike this, both magnetic and electric dipole moments lead to the interconversion of neutrinos of different components. However, in the transversal case of the neutrino polarization, each of these processes can originate through the interaction of charge as well as of any dipole moment with field of emission.

The existence of interconversions $\nu_L \leftrightarrow \nu_R$ and $\bar{\nu}_R \leftrightarrow \bar{\nu}_L$ is incompatible with chiral invariance. These transitions, however, take place owing to the rest mass dependence of the behavior of the neutrino. At our sight, this connection implies that a particle chirality is violated at the expense of the flip of its helicity.

In the case of both longitudinal and transversal polarization of the neutrino, the process (13) is described by the three differential cross sections corresponding to the electric, weak and the united electroweak masses and charges. These cross sections can be defined simultaneously for the same energy if herewith all the three momentum transfer have the space-like size.

One of the beautiful new features of our formulas is the indication to the existence of different low energy limits for the same particle in the interaction type dependence. They of course in the slow neutrino scattering by nuclei behave as

$$E_\nu^E \rightarrow m_\nu^E, \quad E_\nu^{EW} \rightarrow m_\nu^{EW}, \quad E_\nu^W \rightarrow m_\nu^W. \quad (28)$$

At these values the cross sections (22) and (27) describing, respectively, the processes with longitudinal and transversal fermions are not different:

$$d\sigma_{E,W}^{V,A}(\theta_{E,W}, s) = d\sigma_{E,W}^{V,A}(\theta_{E,W}, \varphi, s). \quad (29)$$

For an arbitrary energy such a situation takes place when either vector or axial-vector interactions are present:

$$d\sigma_{E,W}^V(\theta_{E,W}, s) = d\sigma_{E,W}^V(\theta_{E,W}, \varphi, s), \quad (30)$$

$$d\sigma_{E,W}^A(\theta_{E,W}, s) = d\sigma_{E,W}^A(\theta_{E,W}, \varphi, s). \quad (31)$$

But all the three equalities (29)-(31) there exist only at the condition that a particle rest mass does not depend on the type of polarization.

Thus, if it turns out that at the availability of a non-zero mass the longitudinal polarized neutrino must be converted into the transversal polarized and vice versa, this will indicate to the existence of fundamental differences in the masses as well as in the charges of longitudinal and transversal neutrinos.

Comparing (19) with (24), it is easy to observe the contribution of $s(1 - ss')f_{1\nu}^E g_{1\nu}^E$ which is absent at the elastic scattering of longitudinal neutrinos on nuclei but arises as a result of their transversal polarization. The term $s(1 - ss')f_{2\nu}^E g_{2\nu}^E$ available in the longitudinal case of the neutrino polarization does not appear at the transversal particle interaction. We can, therefore, conclude that the invariance of vector and axial-vector types of electroweak currents of longitudinal and transversal neutrinos concerning C, P and T, and also their combinations CP and CPT are different.

Finally, insofar as the spin polarization type dependence of the behavior of massive Majorana neutrinos is concerned, this question together with some aspects of the geometrical nature of inertial mass will be treated in one of our further articles.

References

- [1] E. Fermi, Rend. Lincei **31**, 184, 306 (1922); Phys. Zeit. **23**, 340 (1922).
- [2] R.S. Sharafiddinov, Dokl. Akad. Nauk Ruz. Ser. Math. Tehn. Estest. **7**, 25 (1998).
- [3] R.S. Sharafiddinov, in Proc. Ukrain - Russian Grav. Conf. on Gravitation, Cosmology and Relativistic Astrophysics, Kharkov, November 8-11, 2000 (Kharkov, Ukraine, 2000), p. 25.
- [4] R.S. Sharafiddinov, Spacetime Subst. **1**, 176 (2000).
- [5] R.S. Sharafiddinov, Spacetime Subst. **3**, 47 (2002).
- [6] R.S. Sharafiddinov, Spacetime Subst. **3**, 86 (2002).
- [7] R.S. Sharafiddinov, Spacetime Subst. **3**, 132 (2002).
- [8] M.A.B. Beg, W.J. Marciano and M. Ruderman, Phys. Rev. **D 17**, 1395 (1978).
- [9] W. Bernreuther and M. Suzuki, Rev. Mod. Phys. **63**, 313 (1991).
- [10] R.B. Begzhanov and R.S. Sharafiddinov, Mod. Phys. Lett. **A 15**, (2000) 557; Izv. Russ. Acad. Nauk Ser. Fiz. **64**, 2221 (2000).
- [11] Ya.B. Zel'dovich, Zh. Eksp. Teor. Fiz. **33**, 1531 (1957); Ya.B. Zel'dovich and A.M. Perelomov, Zh. Eksp. Teor. Fiz. **39**, 1115 (1960).
- [12] M.J. Musolf and B.R. Holstein, Phys. Rev. **D 43**, 1956 (1991).
- [13] S. Davidson, B. Campbell and K.D. Bailey, Phys. Rev. **D 43**, 2314 (1991).
- [14] J.A. Morgan and D.B. Farrant, Phys. Lett. **B 128**, 431 (1983).
- [15] T.W. Donnelly and R.D. Peccei, Phys. Rep. **50**, 3 (1979).
- [16] R.S. Sharafiddinov, in Proc. Int. Conf. on Nuclear Physics, St-Peterburg, June 14-17, 2000 (St-Petersburg, 2000), p. 121.